

A New Species of *Selysia* (Cucurbitaceae) from Mesoamerica and a Synopsis of the Genus

Sandra Knapp and Rachel J. Hampshire

Department of Botany, The Natural History Museum, Cromwell Road, London SW7 5BD, United Kingdom

ABSTRACT. A new Panamanian species, *Selysia bidentata*, is described and illustrated, and a key to and a synopsis of *Selysia* are provided. The new species is closely related to *S. smithii*, which occurs from Colombia to Peru in the Andes. This species raises the number of species in *Selysia* to four.

While preparing an account of *Selysia* for *Flora Mesoamericana*, the following new species was identified. It is most similar morphologically to *S. smithii* (Standley) C. Jeffrey from Andean South America and to *S. prunifera* (Poeppig & Endlicher) Cogniaux: differences between the species are detailed in the discussion.

Selysia bidentata R. J. Hampshire, sp. nov. **TYPE:** Panama. Comarca de San Blas: El Llano–Cartí road, 16–20 km from Interamerican Hwy., 9°19'N, 78°55'W, 6 Sep. 1984 (fr), *de Nevers* 3855 (holotype, BM; isotype, MO). Figure 1.

Species *S. pruniferae* (Poeppig & Endlicher) Cogniaux similis, sed seminibus apice bidentatis haud acutis, petiolis villosis nec glabris nec breve pubescentibus.

Monoecious vines; stems sulcate, sparsely pilose to villous; tendrils bifid, often with one branch more robust. Leaves 8–15(–19) × 7–15(–18.5) cm, simple entire and cordate or deeply 3-lobed and broadly ovate to suborbicular, apically acuminate, basally deeply cordate, the margin minutely and remotely denticulate, adaxially glabrate to sparsely tomentose, the veins raised and densely villous, abaxially tomentose; petiole 2.0–2.4(–3.8) cm, loosely villous with straw-colored trichomes 4–5 mm long. Staminate flowers 1–2 in leaf axils; peduncles 3.5–5.2 cm, villous; receptacle conical, villous; calyx 5-lobed, the lobes ca. 6 mm, narrowly triangular; corolla white, campanulate, the lobes ca. 1.3 cm, lanceolate, the margins inrolled, pubescent without, tomentose within. Pistillate flowers 1–2 in leaf axils, only buds seen; ovary ca. 5 mm diam., subglobose, densely tomentose; receptacle cylindrical to subglobose, ca. 3 mm long, calyx lobes 1.5–2 mm, linear; corolla lobes ca. 6 mm in bud. Fruiting peduncle 4.6–8 cm

long, fruit ca. 2.4 cm diam., subglobose, indehiscent, the surface smooth to sparsely short-pubescent, changing from green to red at maturity; seeds 6–9 per fruit, 13–15 × 8–10 mm, erect, ovate, apically 2-dentate, basally 3-dentate, compressed, black or dark brown, the surface minutely pitted.


Selysia bidentata occurs in disturbed forest near roads, streams and trails from 100 to 850 m elevation. A single male flower has been seen (*de Nevers et al.* 5462, MO), but it has not been dissected.

Selysia bidentata is easily distinguished from other species of *Selysia* by its apically bidentate seeds. All other described species in the genus have arrowhead-shaped seeds, with three basal teeth and a single apical point. The epithet “bidentata” refers to this unusual seed shape.

Selysia bidentata is morphologically most similar to *S. smithii* of Andean South America. The two species both have densely villous petioles and sparsely pubescent fruits. Both species also have occasional (in *S. smithii* usually) simple leaves. The petiole pubescence of *S. bidentata* is much longer and looser than that of *S. smithii* and the trichomes are straw-colored rather than ferrugineous. The leaves of *S. smithii* are thick and coriaceous, while those of *S. bidentata* are thinner and more membranous in texture.

The only other Mesoamerican species of *Selysia* is *S. prunifera*, which is easily distinguished from *S. bidentata* by its seed shape and longer, glabrous or minutely pubescent petioles. The petioles of *S. prunifera* exceed 3 cm in length, while those of *S. bidentata* are usually less than 2.4 cm (with rare exceptions).

Paratypes. PANAMA. **Colón:** trail and streamside where Río Escandaloso meets Río Boqueron, 300 ft., 13 Apr. 1978, *Hammel* 2487 (K, MO); along Río Escandaloso and Río Boqueron near their juncture, 28 Apr. 1978, *Hammel* 2740 (MO). **Comarca de San Blas:** Cerro Brewster, premontane rainforest, 850 m, 21 Apr. 1985, *de Nevers et al.* 5462 (MO). **Panamá:** Cerro Jefe to La Eneida, ca. 3,100 ft., 17 Jan. 1968, *Dwyer et al.* 8207 (MO); Cerro Campana, summit and upper trail, 19 Sep. 1975, *Witherspoon & Witherspoon* 8469 (MO).



KEY TO THE SPECIES OF *SELYSIA*

- 1a. Petioles short-pubescent to glabrous, the trichomes to 0.5 mm long.
 - 2a. Leaves with thick tufts of white trichomes in axils of main veins immediately above the petiole insertion point; leaf blades cordate, unlobed 2. *S. cordata*
 - 2b. Leaves without tufts of trichomes in axils of main veins; leaves variously cordate to trilobed 1. *S. prunifera*
- 1b. Petioles villous with long, uniseriate trichomes, the trichomes to 5 mm long.
 - 3a. Petiole trichomes 4–5 mm long, loose, straw-colored; leaves membranous 4. *S. bidentata*
 - 3b. Petiole trichomes 2–3 mm long, very dense, ferrugineous; leaves coriaceous... 3. *S. smithii*

SYNOPSIS OF *SELYSIA* COGNIAUX IN A. DC., MONOGR. PHAN. 3: 735. 1881.

- 1. *Selysia prunifera* (Poeppig & Endlicher) Cogniaux in A. DC., Monogr. Phan. 3: 736. 1881. *Melothria prunifera* Poeppig & Endlicher, Nov. Gen. Sp. Pl. 2: 55, t. 174. 1838. TYPE: Peru. Loreto: Maynas, *Poeppig* 2265 (holotype, W).
Distribution. Costa Rica to Amazonian Peru.

- 2. *Selysia cordata* Cogniaux in A. DC., Monogr. Phan. 3: 736. 1881. TYPE: Colombia. Boyacá: Muzo, *Goudot s.n.* (holotype, P).

Distribution. Northern Colombia in the departments of Santander and Boyacá.

- 3. *Selysia smithii* (Standley) C. Jeffrey, Kew Bull. 25(2): 235. 1971. *Cayaponia smithii* Standley, Publ. Field Mus. Nat. Hist. Bot. Ser. 13(6): 380. 1937. TYPE: Peru. Loreto: Río Morona, *Killip & Smith* 29162 (holotype, US).

Distribution. Andean Colombia to Peru.

- 4. *Selysia bidentata* R. J. Hampshire. TYPE: Panama. Comarca de San Blas: El Llano–Cartí road, *de Nevers* 3855 (holotype, BM).

Distribution. Central Panama.

Acknowledgments. We thank C. Jeffrey (K) for advice, N. K. B. Robson (BM) for help with the Latin diagnosis, and an anonymous reviewer for helpful comments as to relationships in the genus.

←
Figure 1. *Selysia bidentata* R. J. Hampshire. —A. Map of Costa Rica and Panama showing the distribution of *S. bidentata*, solid circles, and *S. prunifera*, open circles. —B. Habit of *S. bidentata*, with female flower and fruit (from *de Nevers* 3855, BM). —C. Male flower (from *de Nevers et al.* 5462, MO). —D. Seed of *S. bidentata* (from *de Nevers* 3855, BM). —E. Seed of *S. prunifera* (from *Kernan* 692, BM).